arXiv:1203.6531v1 [math.GR] 29 Mar 2012

TRANSACTIONS OF THE
AMERICAN MATHEMATICAL SOCIETY
Volume 00, Number 0, Pages 000–000
S 0002-9947(XX)0000-0

GRÖBNER-SHIRSHOV BASIS AND REDUCED WORDS FOR AFFINE WEYL GROUP $\widetilde{A_n}$

EROL YILMAZ, CENAP ÖZEL AND UĞUR USTAOĞLU

Abstract. Using Buchberger-Shirshov Algorithm and Composition-Diamond lemma we obtain the reduced Gröbner-Shirshov bases of $\widetilde{A_n}$ and classify all reduced words of the affine Weyl group $\widetilde{A_n}$.

1. Introduction

Gröbner-Shirshov bases and normal form of the elements were already found for the Coxeter groups of type A_n, B_n and D_n in [1]. They also proposed a conjecture for the general form of Gröbner-Shirshov bases for all Coxeter groups. In [7], an example was given an to show that the conjecture is not true in general. The Gröbner-Shirshov bases of the other finite Coxeter groups are given in [8] and [13]. This paper is another example of finding Gröbner-Shirshov bases for groups, defined by generators and defining relations.

We first cite some concepts and results from the literature which are related to the Gröbner-Shirshov bases for the associative algebras. (see [2, 3, 12])

Suppose S is a linearly ordered set and k is a field. Let $k\langle S\rangle$ be the free associative algebra over k generated by S and S^* be the free monoid generated by S where empty word is the identity which is denoted by 1.

Let S^* be equipped with a monomial ordering $<$. It means that $<$ is a well ordering that agrees with left and right multiplications by words:

$$u > v \Rightarrow w_1uw_2 > w_1vw_2, \text{ for all } w_1, w_2 \in S^*.$$

A standard example of monomial ordering on S^* is deg-lex ordering which first compare two words by length and then comparing them lexicographically where S is a well-ordered set.

Let $f = \alpha\overline{f} + \sum \alpha_i u_i \in k\langle S\rangle$, where $\alpha, \alpha_i \in k, \overline{f} \in S^*$ and $u_i < \overline{f}$ for each i. Then we call $\overline{f}$ the leading word and f monic if $\overline{f}$ has coefficient 1. For a word $w \in S^*$, we denote the length of w by $|w|$.

Definition 1. For two monic polynomial f and g in $k\langle S\rangle$ and a word w, their composition defined by

$$(f,g)_w = \begin{cases} f - agb, & \text{if } w = \overline{f} = a\overline{g}b \\ fb - ag, & \text{if } w = \overline{f}b = a\overline{g}, |\overline{f}| + |\overline{g}| > |w| \end{cases}$$

2010 *Mathematics Subject Classification.* 22E67; 20F55; 51F15; 13P10.

Key words and phrases. Affine Weyl Groups, Gröbner-Shirshov Basis, Composition-Diamond Lemma, q-binomials, Basic Partitions.

©XXXX American Mathematical Society

The word w is called the ambiguity of f and g. The first type of composition is called the composition of including g in f, and the second type is called the composition of intersection of f and g. The transformation $f \mapsto f - agb$ is called the elimination of leading word (ELW) of g in f. Let $R \subset k\langle S\rangle$ be a monic set. The composition $(f,g)_w$ is called reduced to r relative to R if $(f,g)_w = \sum \alpha_i a_i r_i b_i + r$ where every $\alpha_i \in k, a_i, b_i \in S^*, r_i \in R$ with $a_i\overline{r_i}b_i < w$ and the composition $(f,g)_w$ is called trivial relative to R if $r = 0$.

The set $R \subset k\langle S\rangle$ is called Gröbner-Shirshov bases if any composition of polynomials from R is trivial relative to R.

The following lemma was first proved by Shirshov [12] for Lie algebras presented by generators and defining relations. He called it the composition lemma. Similar lemma for free associative algebras was formulated later by Bokut [1] and by Bergman [4] under the name "Diamond lemma" after celebrated Newman's Diamond lemma [9] for graphs. This kind of lemmas are now named as composition-diamond lemmas. We will use Bokut's version of this lemma for free associative algebras. Similar ideas were independently discovered by Hironaka [10] for power series algebras and by Buchberger [5, 6] for polynomial algebras.

Lemma 2. *(Composition-Diamond Lemma for associative algebras)*

Let k be a field, $A = k\langle S|R\rangle = k\langle S\rangle/Id(R)$ and $<$ a monomial ordering on S^*, where $Id(R)$ is the ideal of $k\langle S\rangle$ generated by R. Then the following statements are equivalent:

(i) R is a Gröbner-Shirshov basis.

(ii) $f \in Id(R) \Rightarrow \overline{f} = a\overline{s}b$ for some $s \in R$ and $a, b \in S^*$.

(iii) The set of R-reduced words

$$Red(R) = \{w \in S^* | w \neq a\overline{s}b, a, b \in S^*, s \in R\}$$

is a k-linear basis for the algebra $A = k\langle S|R\rangle$.

For commutative polynomials, this lemma is known as Buchberger's Theorem for the Gröbner basis (see [5, 6]).

If $R \subset k\langle S\rangle$ is not a Gröbner-Shirshov basis, then we reduce every nontrivial compositions to a polynomial relative to R and add this polynomial to the R. Having repeated this procedure (possibly infinitely many times) we obtain a Gröbner-Shirshov basis R^{comp}. The process is called the Buchberger-Shirshov algorithm.

If the set R consists of semigroup relations (i.e. $u - v$, where $u, v \in S^*$), then each nontrivial composition of polynomials from R has the same semigroup form. Hence, R^{comp} consists of semigroup relations too.

Let $A = smg\langle S|R\rangle$ be a semigroup presentation. Then $R \subset k\langle S\rangle$ and we can obtain the Gröbner-Shirshov bases R^{comp}. The set R^{comp} does not depend on the field k and consists of semigroup relations. We will call R^{comp} the Gröbner-Shirshov bases for the semigroup A.

The main purpose of this paper is to find a Grobner-Shirshov basis and classify all reduced words for the affine Weyl group $\widetilde{A}_n$. The strategy for solving the problem is as follows: Let R be the set of polynomials of the defining relations of $\widetilde{A}_n$. Using Buchberger-Shirshov algorithm we obtain new set R' of polynomials including R. Then, by using the algorithm of elimination of leading words with respect to the polynomials in R', all the words in the group $\widetilde{A}_n$ are reduced to the explicit classes of words. After that, we compute the number of the reduced words with respect to these classes by means of a generating function. This generating function turns out the be same with the well known Poincaré polynomial of the affine Weyl group $\widetilde{A}_n$. Therefore, by the Composition-Diamond Lemma the functions in R' form Gröbner-Shirshov basis for the affine Weyl group $\widetilde{A}_n$. Furthermore, one can easily see that this basis is in fact a reduced Gröbner-Shirshov basis.

The results of this paper were obtained during M.Sc studies of Uğur Ustaoğlu at Abant İzzet Baysal University and are also contained in his thesis [14].

2. Gröbner-Shirshov Basis and Reduced Words

Definition 3. The affine Weyl group $\widetilde{A_n}$ has a presentation with generators $S = \{r_0, r_1, \dots, r_n\}$ and defining relations

$r_i r_i = 1 \quad 0 \le i \le n$

$r_i r_j = r_j r_i \quad 0 \le i < j-1 < n$ and $(i,j) \ne (0,n)$

$r_i r_{i+1} r_i = r_{i+1} r_i r_{i+1} \quad 0 \le i \le n-1$

$r_0 r_n r_0 = r_n r_0 r_n$.

Identifying each relation $u = v$ by a polynomial $u - v$, we define

$f_1^{(i)} = r_i r_i - 1 \quad 0 \le i \le n$

$f_2^{(i,j)} = r_i r_j - r_j r_i \quad 0 \le i < j-1 < n$ and $(i,j) \ne (0,n)$

$f_3^{(i)} = r_i r_{i+1} r_i - r_{i+1} r_i r_{i+1} \quad 0 \le i \le n-1$

$f_4 = r_0 r_n r_0 - r_n r_0 r_n$

Let us define

$$r_{ij} = \begin{cases} r_i r_{i+1} \dots r_j, & i < j; \\ r_i r_{i-1} \dots r_j, & i > j; \\ r_i, & i = j; \\ 1, & i = 1, j = 0; \\ 1, & i = n, j = n+1. \end{cases}$$

Lemma 4. *Let $R = \{f_1, f_2, f_3, f_4\}$. A Grobner-Shirshov Basis of $\widetilde{A_n}$ with respect to deglex order with $r_0 > r_1 > \cdots > r_n$ contains the following polynomials.*

$g_1^{(i,j)} = r_{ij} r_i - r_{i+1} r_{ij} \qquad 0 \le i < j-1 < n$ *with* $(i,j) \ne (0,n)$

$g_2 = r_{0n} r_0 r_n - r_1 r_{0n} r_0$

$g_3^{(j,k)} = r_0 r_{nk} r_j - r_j r_0 r_{nk} \qquad 2 \le j < k-1 < n$

$g_4^{(j)} = r_0 r_{nj} r_{j+1} - r_j r_0 r_{nj} \qquad 2 \le j < n$

$g_5^{(k)} = r_0 r_{nk} r_0 - r_n r_0 r_{nk} \qquad 2 \le k < n$

$g_6^{(k,l)} = r_0 r_{nk} r_{1l} r_{0l} - r_n r_0 r_{nk} r_{1l} r_{0,l-1} \qquad 1 \le l < n,\ 2 \le k \le n$

$g_7^{(k,l)} = r_0 r_{nk} r_{1l} r_0 r_{nk} - r_1 r_0 r_{nk} r_{1l} r_0 r_{n,k+1} \qquad 1 \le l < k-1 < n$

$g_8^{(k,l)} = r_0 r_{nk} r_{1l} r_0 r_{n,k-1} - r_1 r_0 r_{nk} r_{1l} r_0 r_{nk} \quad 3 \le k \le n, \quad k-1 \le l \le n$

$g_9^{(j,k,l)} = r_0 r_{nk} r_{1l} r_0 r_{nj} r_{1l} - r_n r_0 r_{nk} r_{1l} r_0 r_{nj} r_{1,l-1}$
$2 \le k \le n-1, \quad k+1 \le j \le n, \quad 1 \le l \le j-2$

$g_{10}^{(j,k,l)} = r_0 r_{nk} r_{1l} r_0 r_{nj} r_{1,l+1} - r_n r_0 r_{nk} r_{1l} r_0 r_{nj} r_{1l}$
$2 \le k \le n, \quad k \le j \le n, \quad j-1 \le l \le n-1$

Proof. Let $R = \{f_1, f_2, f_3, f_4\}$. We apply the Buchberger-Shirshov algorithm to the R. We show every ELW in the below computations except ELW's of $f_2(i,j)$, the cummutativity relations. Notice that at this point we are not claiming that this is a Gröbner-Shirshov basis for $\widetilde{A}_n$.

$(f_3^{(i)}, f_2^{(i,i+2)}) = g_1^{(i,i+2)}$.

$(g_1^{(i,j-1)}, f_2^{(i,j)}) = g_1^{(i,j)} \quad \text{for} \quad j = i+3, \ldots, n$.

$(g_1^{(0,n-1)}, f_4) = g_2$.

$(f_2^{(0,j)}, f_2^{(j,n)}) = g_3^{(j,n)}$.

$(g_3^{(j,k+1)}, f_2^{(j,k)}) = g_3^{(j,k)} \quad \text{for} \quad k = n-1, \ldots, j+2$.

$(f_2^{(0,n-1)}, f_3^{(n-1)}) = g_4^{(n-1)}$.

$(g_3^{(j,j+2)}, f_3^{(j)}) = g_4^{(j)} \quad \text{for} \quad j = n-2, ..., 2$.

$(f_4, f_2^{(0,n-1)}) = g_5^{(n-1)}$.

$(g_5^{(k+1)}, f_2^{(0,k)}) = g_5^{(k)} \quad \text{for} \quad k = n-2, \ldots, 2$.

$(f_4, f_3^{(0)}) = g_6^{(n,1)}$

$(g_5^{(k)}, f_3^{(0)}) = g_6^{(k,1)} \quad \text{for} \quad k = n-1, \ldots, 2$.

$(g_6^{(k,l-1)}, f_3^{(l-1)}) = g_6^{(k,l)} \quad \text{for} \quad l = 2, \ldots, n-1$.

$(f_3^{(0)}, f_4) = r_0 r_n r_1 r_0 r_n - r_1 r_0 r_n r_1 r_0 = g_7^{(n,1)}$.

$(g_1^{(0,l)}, f_4) = g_7^{(n,l)} \quad \text{for} \quad l = 2, \ldots, k-2$.

$$\begin{aligned}(g_7^{(k+1,l)}, g_3^{(k+1,k)}) &= r_0 r_{n,k+1} r_{1l} r_k r_0 r_{nk} - r_1 r_0 r_{n,k+1} r_{1l} r_0 r_{n,k+2} r_k r_{k+1} \\ &= g_7^{(k,l)} - r_1 r_0 r_{n,k+1} r_{1l} g_3^{(k,k+2)} r_{k+1} \quad \text{for} \quad k = n-2, \ldots, 3.\end{aligned}$$

$$\begin{aligned}(g_7^{(n,l)}, g_3^{(n,n-1)}) &= r_0 r_n r_{1l} r_n r_0 r_{n,n-1} - r_1 r_0 r_n r_{1l} r_0 r_{n,n-1} r_n \\ &= g_7^{(n-1,l)} - r_1 r_0 r_n r_{1l} f_2^{(0,n-1)} r_n \quad \text{for} \quad k = n-1.\end{aligned}$$

$$\begin{aligned}(g_7^{(k,k-2)}, g_4^{(k-1)}) &= r_0 r_{nk} r_{1,k-1} r_0 r_{n,k-1} - r_1 r_0 r_{nk} r_{1,k-2} r_0 r_{n,k+1} r_{k-1} r_k \\ &= g_8^{(k,k-1)} - r_1 r_0 r_{nk} r_{1,k-2} g_3^{(k-1,k+1)} r_n \quad \text{for} \quad k = n-1, \ldots, 3.\end{aligned}$$

$$\begin{aligned}(g_7^{(n,n-2)}, g_4^{(n-1)}) &= r_0 r_n r_{1,n-1} r_0 r_{n,n-1} - r_1 r_0 r_n r_{1,n-2} r_0 r_{n-1} r_n \\ &= g_8^{(n,n-1)} - r_1 r_0 r_n r_{1,n-2} f_2^{(0,n-1)} r_n \quad \text{for} \quad k = n.\end{aligned}$$

$$(g_2, g_4^{(n-1)}) = r_{0,n-2} f_3^{(n-1)} r_0 r_n r_{n-1} - r_1 r_{0n} f_2^{(0,n-1)} r_n - r_1 r_{0,n-2} f_3^{(n-1)} r_0 r_n + g_8^{(n,n)}$$

$$\begin{aligned}(g_8^{(k+1,l)}, g_4^{(k-1)}) &= r_0 r_{n,k+1} r_{1,k-2} g_1^{(k-1,l)} r_0 r_{n,k-1} - r_1 r_0 r_{n,k+1} r_{1l} g_3^{(k-1,k+1)} r_k \\ &\quad - r_1 r_0 r_{n,k+1} r_{1,k-2} g_1^{(k-1,l)} r_0 r_{n,k+1} r_k + g_8^{(k,l)} \quad \text{for} \quad l = k, \ldots, n.\end{aligned}$$

$(g_6^{(k,1)}, f_2^{(1,n)}) = g_9^{(k,n,1)}$.

$$(g_9^{(k,j+1,1)}, f_2^{(1,j)}) = g_9^{(k,j,1)} \quad \text{for} \quad j = n-1, \dots, k+1.$$

$$\begin{aligned}(g_9^{(k,j,l-1)}, f_3^{(l-1)}) &= r_0 r_{nk} r_{1,l-1} r_0 r_{nj} r_{1,l-2} r_l r_{l-1} r_l - r_n r_0 r_{nk} r_{1,l-1} r_0 r_{nj} r_{1,l-2} r_l r_{l-1} \\ &= g_9^{(k,j,l)} \quad \text{for} \quad l = 2, \dots, j-2.\end{aligned}$$

$$\begin{aligned}(g_9^{(k,j+1,j-1)}, f_3^{(j-1)}) =& r_0 r_{nk} r_{1,j-1} r_0 r_{n,j+1} r_{1,j-2} r_j r_{j-1} r_j \\ &- r_n r_0 r_{nk} r_{1,j-1} r_0 r_{n,j+1} r_{1,j-2} r_j r_{j-1} \\ =& g_{10}^{(k,j,j-1)} \quad \text{for} \quad j = n-1, \dots, k.\end{aligned}$$

$$\begin{aligned}(g_6^{(k,n-1)}, f_3^{(n-1)}) =& r_0 r_{nk} r_{1,n-1} r_{0,n-2} r_n r_{n-1} r_n \\ &- r_n r_0 r_{nk} r_{1,n-1} r_0 r_n r_{1,n-1} = g_{10}^{(k,n,n-1)}.\end{aligned}$$

$$\begin{aligned}(g_{10}^{(k,j,l-1)}, f_3^{(l)}) =& r_0 r_{nk} r_{1,l-1} r_0 r_{nj} r_{1,l-1} r_{l+1} r_l r_{l+1} \\ &- r_n r_0 r_{nk} r_{1,l-1} r_0 r_{nj} r_{1,l-1} r_l r_{l+1} r_l \\ =& r_0 r_{nk} r_{1,l-1} g_4^{(l)} r_{l-1,j} r_{1,l+1} - r_n r_0 r_{nk} r_{1,l-1} g_4^{(l)} r_{l-1,j} r_{1l} \\ &+ g_{10}^{(k,j,l)} \quad \text{for} \quad l = j-1, \dots, n-1.\end{aligned}$$

In the above equation if $l = j$, then $r_{l-1,j}$ assumed to be the identity 1.

□

Let $R' = R \cup \{g_1, \dots, g_{10}\}$. We want to find the properties of the elements of the set $Red(R') = \{w \in S^* | w \neq a\bar{f}b, a, b \in S^*, f \in R'\}$. If $w \in Red(R')$, then we call it a reduced word.

Notice that elements of R' not containing r_0 is in fact a Gröbner-Shirshov basis for the Coxeter group A_n. The following lemma is just another way of expressing of the Lemma 3.2 of [1].

Lemma 5. *Any reduced word not containing r_0 is in the form*

$$r = (r_{nj_n})^{\alpha_n} (r_{n-1,j_{n-1}})^{\alpha_{n-1}}, \dots, (r_{2j_2})^{\alpha_2}, (r_{1j_1})^{\alpha_1}$$

where $i \leq j_i \leq n$ and $\alpha_i \in \{0, 1\}$.

After investigation of leading words of the elements of G', we can claim the following results. For convenience we write $r_0 r_{n,n+1} r_{1l}$ instead of r_{0l} and $r_0 r_{nk} r_{10}$ instead of $r_0 r_{nk}$.

Lemma 6. *The following words are reduced.*

(i) $w = r_0 r_{nk} r_{1l}$ $2 \leq k \leq n+1$, $0 \leq l \leq n$

(ii)

$$(r_0 r_{nk} r_{1l})(r_0 r_{np} r_{1q}) = \begin{cases} (k < p) \wedge (l > q), & \textit{if} \quad q - p < l - k < -1 \\ (k \leq p) \wedge (l > q), & \textit{if} \quad (q - p < -1) \wedge (l - k \geq -1) \\ (k \leq p) \wedge (l \geq q), & \textit{if} \quad l - k > q - p \geq -1 \end{cases}$$

Lemma 7. *Let w be a reduced word starting with r_i for $i = 1, \dots, n$ and let t be a reduced word starting with r_0. Then wt is also a reduced word.*

Proof. The results follows from the following observation. Any leading word starting with r_i for $i = 1, \dots, n$ in G' do not contains r_0. ☐

Lemma 8. *Let $w_1, w_2, \dots w_k, w_{k+1}$ be reduced words in the one of the forms given in the first three items of Lemma 6. If $w_1 w_2 \dots w_k$ and $w_2 w_3 \dots w_{k+1}$ are reduced, then $w_1 w_2 \dots w_{k+1}$ is also reduced.*

Proof. Let $w = w_1 w_2 \dots w_{k+1}$. The only possible reduced subword of w is in the form $r w_2 w_3 \dots w_k s$ where r and s are subwords of w_1 and w_{k+1}, respectively. Since $w_2 w_3 \dots w_k w_{k+1}$ is reduced word, $w_2 w_3 \dots w_k s$ is also reduced word starting with r_0. Since r is a subword of w_1, r is also a reduced word not containing r_0. By Lemma 7, $r w_2 w_3 \dots w_k s$ is also reduced. Hence $w_1 w_2 \dots w_{k+1}$ must be a reduced word. ☐

Definition 9. Let $a_i = r_0 r_{nk} r_{1l}$ for $2 \leq k \leq n+1$, $0 \leq l \leq n$ and $l - k = i - 2$.

Let

$$u = (a_n)^{m_n} (a_{n-1})^{m_{n-1}} \dots (a_1)^{m_1}$$

where $m_i \geq 0$ for $i = 1, \dots, n$. Furthermore, if $a_i = r_0 r_{nk} r_{1l}$, then $a_{i+1} = r_0 r_{n,k+1} r_{1l}$ or $a_{i-1} = r_0 r_{nk} r_{1,l-1}$. Notice that the number of possible u's is 2^{n-1}. We call a_i's the components of the word u.

Definition 10. Let $b_t = r_0 r_{np} r_{1q}$ for $2 \leq p \leq n+1$ and $0 \leq q \leq n$ satisfying $q - p < -1$. There are $n!$ such words.

Let

$$v^{(b_t)} = b_t (b_{t-1})^{\alpha_{t-1}} \cdots (b_s)^{\alpha_s}$$

where $\alpha_i \in \{0, 1\}$. Furthermore, if $b_i = r_0 r_{np_i} r_{1q_i}$, then $b_{i-1} = r_0 r_{np_{i-1}} r_{1q_{i-1}}$ for $p_i < p_{i-1}$ and $q_i > q_{i-1}$. If $p_i = n+1$ or $q_i = 0$, then $\alpha_j = 0$ for $j = s, \dots, i-1$.

For the convenience, we define $1 = r_0 r_{n,\infty} r_{1,-1}$ and $v(1) = 1$.

Proposition 11. *Let u and $v^{(b_t)}$ be words defined above where $a_1 = r_0 r_{nk} r_{1l}$ and $b_t = r_0 r_{np} r_{1q}$. Then the words u, $v^{(b_t)}$ and $w = uv^{(b_t)}$ are reduced if $p \geq k$ and $q < l$ in w.*

Proof. The result is easily follows from Lemma 6 and Lemma 8. ☐

Figure 1.1 shows every possible reduced words for $n = 4$.

Let

$$w_1 = (r_0 r_{42} r_{14})^{m_1} (r_0 r_{42} r_{13})^{m_2} (r_0 r_{43} r_{13})^{m_3} (r_0 r_4 r_{13})^{m_4} r_{01}$$

and

$$w_2 = (r_0 r_{42} r_{14})^{m_1} (r_0 r_{43} r_{14})^{m_2} (r_0 r_4 r_{14})^{m_3} (r_0 r_4 r_{13})^{m_4} r_{01}.$$

Then w_1 and w_2 are two reduced words in $\widetilde{A}_4$. If we take $m_2 = m_3 = 0$ and $m_1 = m_4 = 1$, then the subword $(r_0 r_{42} r_{14})(r_0 r_4 r_{13}) r_{01}$ is written twice. To avoid this situation, we define the arranged words.

Definition 12. Let $w = uv^{(b_t)}$ be a reduced word where $b_t = r_0 r_{np} r_{1q}$ and

$$u = (a_n)^{(m_n)} (a_{n-1})^{(m_{n-1})} \cdots (a_1)^{(m_1)}.$$

For $i = 2, \dots, n-1$, let $m_i \geq 1$ if $a_{i+1} = r_0 r_{nk} r_{1,l+1}$, $a_i = r_0 r_{nk} r_{1l}$ and $a_{i-1} = r_0 r_{n,k+1} r_{1l}$. If $a_2 = r_0 r_{nk} r_{1,l+1}, a_1 = r_0 r_{nk} r_{1l}$ and $p > k$, then let $m_1 \geq 1$.



FIGURE 1. 1

Then w is called an arranged word and the components a_i where $m_i \geq 1$ are called a marked component of w.

Theorem 13. *If all reduced words $w = uv^{(b_t)}$ are arranged, then each subword is written uniquely.*

Proof. Let $w_1 = u_1v^{(b_t)}$ and $w_2 = u_2v^{(b_t)}$ be two arranged word. Since the only possibility for $a_n = r_0r_{n2}r_{1n}$, the first components of them are the same. Let us assume that $a_n, a_{n-1}, \dots, a_j$ are common in w_1 and w_2. If the exponents of a_i for $i = j, \dots, n$ were 1 and the others were 0 in both w_1 and w_2, then the word $a_na_{n-1}\cdots a_j$ would be written twice.

Let us assume $a_j = r_0r_{nk}r_{1l}$. Moreover, let $a_{j-1} = r_0r_{n,k+1}r_{1l}$ in w_1 and $a_{j-1} = r_0r_{nk}r_{1,l-1}$ in w_2. Then there exits $s \geq l-1$ the components of w_2 between $j-1$ and $s+1$ are

$$(r_0r_{nk}r_{1,l-1})(r_0r_{nk}r_{1,l-2})\cdots(r_0r_{nk}r_{1s})(r_0r_{n,k+1}r_{1s})$$

when $s-k > -1$. If $s-k = -1$, we have

$$(r_0r_{nk}r_{1,l-1})(r_0r_{nk}r_{1,l-2})\cdots(r_0r_{nk}r_{1s})(r_0r_{np}r_{1q})$$

Therefore $r_0r_{nk}r_{1s}$ is a marked component in the first case. By Definition 9, $r_0r_{nk}r_{1l}$ can not be a component of w_1. The component of w_1 in the same position can be $r_0r_{n\overline{k}}r_{1\overline{s}}$ where $\overline{k} > k$ and $\overline{s} < s$. Since b_t is common in both words, $p \geq \overline{k} > k$. Therefore $r_0r_{nk}r_{1s}$ is also a marked component in the second case.Hence the word $(a_n)a_{n-1}\cdots(a_j)$ can be written only in w_1 not in w_2.

□

3. COUNTING REDUCED WORDS

Theorem 14. *Let $\widehat{w} = a_{i_s}a_{i_{s-1}}\dots a_{i_1}v^{(b_t)}$ where $a_{i_j} = r_0r_{nk_j}r_{nl_j}$.Then a_{i_j}'s are marked components of an arranged word $w = uv^{(b_t)}$ where $b_t = r_0r_{np}r_{1q}$ if and only if $k_s < k_{s-1} < \dots < k_1 < p$ and $q < l_1 < l_2 < \dots < l_s < n$*

Proof. By construction of arranged words, the marked components satisfy the given conditions. Conversely, if $\widehat{w} = a_{i_s} a_{i_{s-1}} \dots a_{i_1} v^{(b_t)}$ is a word satisfying the conditions, then an arranged word ($w = uv^{(b_t)}$) whose marked components are a_{i_j}'s can be obtained as follows:

The components up to $r_0 r_{nk_s} r_{1l_s}$ are

$$(r_0 r_{n2} r_{1n}) \cdots (r_0 r_{nk_s} r_{1n})(r_0 r_{nk_s} r_{1,n-1}) \cdots (r_0 r_{nk_s} r_{1l_s}),$$

the components between $(r_0 r_{nk_j} r_{1l_j})$ and $(r_0 r_{nk_{j-1}} r_{1l_{j-1}})$ are

$$(r_0 r_{nk_j} r_{1l_j}) \cdots (r_0 r_{nk_{j-1} r_{1l_j}})(r_0 r_{nk_{j-1}} r_{1,l_j-1}) \cdots (r_0 r_{nk_{j-1}} r_{1l_{j-1}})$$

and the last part of the word$w = uv^{(b_t)}$ is

$$(r_0 r_{nk_2} r_{1l_2}) \cdots (r_0 r_{nk_1 r_{1l_2}})(r_0 r_{nk_1} r_{1,l_2-1}) \cdots (r_0 r_{nk_1} r_{1l_1})(v^{(b_t)}).$$

□

Therefore the number of the elements in an arranged word $w = uv^{(b_t)}$ given by the generating function

$$\frac{x^{\alpha}}{(1-x^{2n})(1-x^{2n-1})\cdots(1-x^{n+1})}$$

where α is the length of the word $\widehat{w}$. In order to count all reduced words starting with r_0 we have to find the number words $\widehat{w}$ whose length is α for any power α. To do this, we will find a correspondence between these words and some special partitions of integers.

If m is a positive integer, then a partition of m is a nonincreasing sequence of positive integers $p_1, p_2, \dots, p_k$ whose sum is m. Each p_i is called a part of the partition. Let n be a positive integer. Any partition $m = d_1 + d_2 + \dots + d_k$ where $k \leq n$ can be identify by the $n-$tuple $(d_1, d_2, \dots, d_k, 0, 0, \dots, 0)$.

We can also represent each word $r_0 r_{nk} r_{1l}$ with the n-tuple $(k, 1, \dots, 1, 0, \dots, 0)$, r_{nk} with $(k, \dots, 0)$ and r_{0l} with $(1, 1 \dots, 1, 0, \dots, 0)$ where number one 1's is equal to l for $2 \leq k \leq n-1, 1 \leq l \leq n$.

Definition 15. Let n be a positive integer. The n-tuples $(k, 1 \dots, 1, 0, \dots, 0)$ where number of 1's is l for $1 \leq k \leq n$, $1 \leq l \leq n-1$ are called basic partitions. The basic partition $(k_1, 1 \dots, 1, 0 \dots, 0)$ is said to be connected to the basic partition $(k_2, 1, \dots, 1, 0, \dots, 0)$ if $k_1 > k_2$ and the number of 1's in the first one is greater than number of 1's in the second one. Hence a sequence of connected partition $a_1, a_2, \dots, a_m$ corresponds to a word $\widehat{w}$ given in Theorem 14.

Theorem 16. *There is one to one correspondence between words $\widehat{w}$ and the partitions in which there are at most n parts and in which no parts is larger than n.*

Proof. Since we identify each word $\widehat{w}$ with a sequence of connected basic partitions, we must find a correspondence between sequences of connected partitions and the partitions fit into a box of size $n \times n$. Let $a_1, a_2, \dots, a_m$ be a sequence of connected partitions where $a_i = (k_i, \underbrace{1, \dots, 1}_{l_i} 0, \dots, 0)$. Hence $k_i > k_j$ and $l_i > l_j$ for $1 \leq i < j \leq m$.

Define

$$\bigoplus_{i=1}^{m} a_i = \sum_{i=1}^{m} \sigma_1^{i-1}(a_i)$$

where $\sigma(p_1, p_2, \ldots, p_{n-1}, p_n) = (p_n, p_1, p_2, \ldots, p_{n-1})$. Then

$$\bigoplus_{i=1}^{m} a_i = (k_1, k_2{+}1, \ldots, k_m{+}m{-}1, \underbrace{m, \ldots, m}_{l_m}, \underbrace{m-1, \ldots, m-1}_{l_{m-1}-(l_m+1)}, \ldots, \underbrace{1, \ldots, 1}_{l_1-(l_2+1)}, 0 \ldots, 0)$$

We prove the last equation by induction on m.

Let $m = 2$, Since $l_1 - 1 \geq l_2$,

$$\begin{aligned} a_1 \oplus a_2 &= (k_1, \underbrace{1, \ldots, 1}_{l_1}, 0, \ldots, 0) + (0, k_2, \underbrace{1, \ldots, 1}_{l_2}, 0, \ldots, 0) \\ &= (k_1, k_2 + 1, \underbrace{2, \ldots, 2}_{l_2}, \underbrace{1, \ldots, 1}_{l_1-(l_2+1)}, 0, \ldots, 0) \end{aligned}$$

Let us assume that

$$\bigoplus_{i=1}^{m-1} a_i = (k_1, k_2 + 1, \ldots, k_{m-1} + m - 2, \underbrace{m-1, \ldots, m-1}_{l_{m-1}}, \ldots, \underbrace{1, \ldots, 1}_{l_1-(l_2+1)}, 0, \ldots, 0).$$

$$\begin{aligned} \bigoplus_{i=1}^{m} a_i &= \sum_{i=1}^{m} \sigma^{i-1}(a_i) \\ &= \sum_{i=1}^{m-1} \sigma^{i-1}(a_i) + \sigma^{m-1}(a_m) \\ &= \bigoplus_{i=1}^{m-1} a_i + \sigma^{m-1}(a_m) \\ &= (k_1, k_2 + 1, \ldots, k_{m-1} + m - 2, \underbrace{m-1, \ldots, m-1}_{l_{m-1}}, \ldots, \underbrace{1, \ldots, 1}_{l_1}, 0, \ldots, 0) \\ &\quad + (\underbrace{0, 0, \ldots, 0}_{m-1}, k_m, \underbrace{1, \ldots, 1}_{l_m}, 0, \ldots, 0) \\ &= (k_1, k_2 + 1, \ldots, k_m + m - 1, \underbrace{m, \ldots, m}_{l_m}, \ldots, \underbrace{1, \ldots, 1}_{l_1-(l_2+1)}, 0, \ldots, 0) \end{aligned}$$

The last equality easily follows from the fact $l_{m-1} - 1 \geq l_m$. Since $n \geq k_1 \geq k_2 + 1 \geq \cdots > k_m + (m-1) \geq m$, the last line corresponds to a partition of $2n$ fits into a n by n box.

Conversely let $m = (m_1, m_2, \ldots, m_n)$ be a partition where $n \geq m_1 \geq m_2 \geq \cdots \geq m_n \geq 0$. If i_1 is the last index such that $m_{i_1} \neq 0$, then let $a_1 = (m_1, 1, \ldots, 1, 0, \ldots 0)$ where the last 1 in i_1-th position. Then let

$$x = \sigma^{-1}(m - a_1) = (m_2 - 1, \ldots, m_{i_1} - 1, 0, \ldots, 0)$$

and $a_2 = (m_2 - 1, \ldots, 1, 0, \ldots, 0)$ where the position of the last nonzero element in x and the position of last 1 in a_2 are same. Clearly, a_1 and a_2 are basic partitions and a_1 is connected to a_2. Continuing the process until reaching the $(0, \ldots, 0)$, one can obtain a sequence of connected basic partitions. $\square$

Definition 17. Let be positive integers. The q-binomial is defined by

$$\left(\begin{array}{c} m \\ r \end{array} \right)_q = \frac{(1-q^m)(1-q^{m-1})\cdots(1-q^{m-r+1})}{(1-q)(1-q^2)\cdots(1-q^r)}$$

Although the formula in the first clause appears to involve a rational function, it actually designates a polynomial, because the division is exact in $\mathbb{Z}[q]$. A standard combinatorial interpretation for q-binomial is that it counts the number of partitions that will fit into a box of size $k \times (n-k)$, weighted by the size of the partition. In particular the q-binomial

$$\left(\begin{array}{c} 2n \\ n \end{array} \right)_x = \frac{(1-x^{2n})(1-x^{2n-1})\cdots(1-x^{n+1})}{(1-x)(1-x^2)\cdots(1-x^n)}$$

counts the number of partitions in which there are at most n parts and in which no parts is larger than n.

Now, we can proof the main result of this paper.

Theorem 18. *Then the reduced Gröbner-Shirshov basis of the affine Weyl group $\widetilde{A}_n$ is the set G'. Moreover all the reduced words are the form rw where r is a reduced word not including r_0 and w is a arranged word.*

The reduced words not including r_0 is given in Lemma 5. It is easy to see that the number of such words given by the generating function

$$(1+x)(1+x+x^2)\cdots(1+x+\cdots+x^n).$$

Theorem 14 and Theorem 16 imply that the number of arranged words given by the generating function

$$\begin{aligned} \frac{\left(\begin{array}{c} 2n \\ n \end{array} \right)_x}{(1-x^{2n})(1-x^{2n-1})\cdots(1-x^{n+1})} &= \frac{\frac{(1-x^{2n})(1-x^{2n-1})\cdots(1-x^{n+1})}{(1-x)(1-x^2)\cdots(1-x^n)}}{(1-x^{2n})(1-x^{2n-1})\cdots(1-x^{n+1})} \\ &= \frac{1}{(1-x)(1-x^2)\cdots(1-x^n)}. \end{aligned}$$

By Lemma 7, the reduced words of $\widetilde{A}_n$ are in the form rw. Hence the number of reduced words given by the generating function

$$\frac{(1+x)(1+x+x^2)\cdots(1+x+\cdots+x^n)}{(1-x)(1-x^2)\cdots(1-x^n)}$$

which is well known Poincaré polynomial of the affine Weyl group $\widetilde{A}_n$. (see [11]). Therefore these are all reduced words of $\widetilde{A}_n$. Hence by Composition-Diamond Lemma, G' is a Gröbner-Shirshov basis of $\widetilde{A}_n$. In fact, G' is a reduced Gröbner-Shirshov basis.

REFERENCES

[1] L.A. Bokut and L.S. Shiao, *Gröbner-Shirshov bases for Coxeter groups*, Comm. Algebra **29** (2001), 4305–4319.

[2] L. A. Bokut, *Imbeddings into Simple Associative Algebras*, Algebra Logika **15** (1976), 117-142.

[3] L.A. Bokut, *Unsolvability of the word problem, and subalgebras of finitely presented Lie algebras*, Izv. Akad. Nauk. SSSR Ser. Mat., **36** (1972), 1173-1219.

[4] G. M. Bergman,*The Diamond Lemma for Ring Theory*, Adv. Math. **29** (1978), 178-218.

[5] B. Buchberger, *An Algorithm for Finding a Basis for the Residue Class Ring of a Zero-dimensional Polynomial Ideal*, Ph.D. Thesis, University of Innsbruck, Austria, 1965.

[6] B. Buchberger, *An Algorithmical Criteria for the Solvability of Algebraic Systems of Equations*, Aequationes Math. 4 (1970), 374-383.

[7] Y.Chen and C. Liu, *Gröbner-Shirshov bases for Coxeter groups I*, arXiv:0910.0096v1.

[8] D. Lee, *Gröbner-Shirshov bases and normal forms for the coxeter groups E_6 and E_7*, Advances in Algebra and Combinatorics, World Scientific, 2008, 243–255.

[9] M. H. A. Newman, *On Theories with a Combinatorial Definition of Equivalence*, Ann. Math. **43** (1942), 223-243.

[10] H. Hironaka, *Resolution of Singulatities of an Algebtaic Variety over a Field of Characteristic Zero, I, II*, Ann. Math. **79** (1964), 109-203, 205-326.

[11] J. E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge Studies in Advanced Mathematics, **29**, Cambridge University Press, Cambridge, (1990).

[12] A. I. Shirshov, *Some Algorithmic Problem for Lie Algebras*, Sib. Mat. Zh. **3** (1962), 292-296. English translation: SIGSAM Bull. **33 (2)**, (1999), 3-6.

[13] O. Svechkarenko, *Gröbner-Shirshov bases for the Coxeter group E_8*, Master Thesis, Novosibirsk State University, (2007).

[14] U. Ustaoğlu, *Gröbner-Shirshov bases and reduced forms for the affine Weyl groups*, Master Thesis, Abant İzzet Baysal University, (2012).

Department of Mathematics, Abant İzzet Baysal University, Bolu, Turkey
E-mail address: `yilmaz_e2@ibu.edu.tr, cenap@ibu.edu.tr, ugur1987@gmail.com`